

On the nature of "hidden symmetry" or "accidental degeneracy" in the Kepler problem

Tamar T. Khachidze*, Anzor A. Khelashvili*

* *Department of Theoretical Physics, Ivane Javakhishvili Tbilisi State University,
I.Chavchavadze ave. 3, 0128, Tbilisi, Georgia*

February 2, 2008

Lecture given at the *Summer School and Conference "New Trends of High Energy Physics"*,
Yalta, Crimea, Ukraine, 16-23 Sept. 2006

Abstract

We propose a symmetry of the Dirac equation under the interchange of signs of eigenvalues of the Dirac's K operator. We show that the only potential which obeys this requirement is the Coulomb one for both vector and scalar cases.

The problem is rather old, but several principal questions remain without response. The Kepler problem has an additional symmetry, more precisely, additional conserved quantity, so-called Laplace-Runge-Lenz (LRL) vector [1]. Because of this fact there appears extended algebra together with orbital momentum, which is isomorphic to $SO(4)$ (for negative total energies). But components of LRL vector as generators of this algebra do not participate in any geometric transformations. In 1926 W.Pauli [2] considered this algebra in quantum mechanics and obtained Hydrogen atom spectrum by only algebraic methods. In 30-ies V.Fock [3] considered the Schroedinger equation for the Kepler problem in momentum representation and had shown that the Coulomb spectrum has an $SO(4)$ symmetry in energy-momentum space. Afterwards algebraic methods attract more wide applications and it was cleared up in the last decades that the hidden symmetry of the Kepler problem is closely related to the supersymmetry of Hydrogen atom.

Because of its significance this symmetry appears even in the textbooks. For example, in Classical Mechanics by J.Goldstein [4], as well as in recent books of Quantum Mechanics [5]. As regards of relativistic quantum mechanics (Dirac equation) relatively less is known, although the so-called radial supersymmetry was demonstrated a long-time ago [6] and the many problems are studied rather well.

The aim of our talk is to determine what the Dirac equation tells us about this problem. We can see that very interesting physical picture emerges.

For the Kepler problem in the Dirac equation Johnson and Lippmann [7] published very brief abstract in Physical Review at 1950. They had written that there is an additional conserved quantity

$$A = \vec{\sigma} \cdot \vec{r} r^{-1} - i \left(\frac{\hbar c}{e^2} \right) (mc^2)^{-1} j \rho_1 (H - mc^2 \rho_3), \quad (1)$$

which plays the same role in Dirac equation, as the LRL vector in Schroedinger equation.

As regards to the more detailed derivation, to the best our knowledge, is not published in scientific literature (one of the curious fact in the history of physics of 20th century) Moreover as far as commutativity of the Johnson-Lippmann(JL) operator with the Dirac Hamiltonian is concerned, it is usually mentioned that it can be proved by "rather tedious calculations" [8].

Recently [9, 13], we developed rather simple and transparent way for deriving the JL operator. We obtained this operator and at the same time proved its commutativity with the Hamiltonian.

After we consider necessity to be convinced, that the Coulomb problem is distinguishable in this point of view. We considered the Dirac equation in arbitrary central potential, $V(r)$ and shown that the symmetry, which will be defined more precisely below, takes place only for Coulomb potential.

We have chosen the subject of today's talk so that it is consonant with 1966 School, 40th Years of which today's School is devoted to, when several lectures were dedicated to the problem of hidden symmetry - lectures by Alilluev, Matveenkov, Popov and Werle [10]. During several decades these lectures were the only source, from which the young students and scientists could learn about this topic and the Proceedings of that School is an excellent monograph in many subjects of Theoretical High Energy Physics.

In some respects our today's lecture is a continuation (but not analytic) of those old lectures.

First of all consider the Dirac Hamiltonian in central field, $V(r)$, which is the 4th component of the Lorentz vector:

$$H = \vec{\alpha} \cdot \vec{p} + \beta m + V(r) \quad (2)$$

The so-called Dirac's K -operator, defined as

$$K = \beta(\vec{\Sigma} \cdot \vec{l} + 1), \quad (3)$$

commutes with this Hamiltonian for arbitrary $V(r)$, $[K, H] = 0$. Here $\vec{\Sigma}$ is the spin matrix

$$\vec{\Sigma} = \begin{pmatrix} \vec{\sigma} & 0 \\ 0 & \vec{\sigma} \end{pmatrix}$$

and $\vec{l}$ - orbital momentum vector-operator.

It is evident that the eigenvalues of Hamiltonian also are labeled by eigenvalues of K . For example, the well-known **Sommerfeld formula** for the Coulombic spectrum looks like

$$E_{n,|\kappa|} = m \left[1 + \frac{(Z\alpha)^2}{(n - |\kappa| + \sqrt{\kappa^2 - (Z\alpha)^2})^2} \right]^{-1/2}$$

It manifests explicit dependence on κ , more precisely on $|\kappa| = j + 1/2$. It is surprising that for other solvable potentials the degeneracy with respect to signs of κ does not take place.

Is it peculiar only for Coulomb potential?

Let us consider Hamiltonian (2) with arbitrary potential and **require the degeneracy in $\pm\kappa$** . It is natural that for description of this degeneracy one has to find an operator, that mixes these two signs. It is clear that such an operator, say A , must be anticommuting with K , i.e.

$$\{A, K\} = AK + KA = 0 \quad (4)$$

If at the same time this operator commutes with the Hamiltonian, then it'll generate the symmetry of the Dirac equation. Therefore, we need an operator A with the following properties:

$$\{A, K\} = 0 \quad \text{and} \quad [A, H] = 0 \quad (5)$$

It is our definition of symmetry we are looked for.

It is interesting that after this operator is constructed, we will be able to define relativistic supercharges as follows:

$$Q_1 = A, \quad Q_2 = i \frac{AK}{|\kappa|}$$

It is obvious that

$$\{Q_1, Q_2\} = 0, \quad Q_1^2 = Q_2^2$$

and we can find Wittens superalgebra [12], where $Q_1^2 \equiv h$ is a so-called, Witten's Hamiltonian.

Now our goal is a construction of the operator A . We know, that there is a Diracs γ^5 matrix, that anticommutes with K . **What else?** There is a **simple theorem** [13], according to which arbitrary $(\vec{\Sigma} \cdot \vec{V})$ **type operator, where $\vec{V}$ is a vector with respect to $\vec{l}$ and is perpendicular to it, $(\vec{l} \cdot \vec{V}) = (\vec{V} \cdot \vec{l})$, anticommutes with K :**

$$\{(\vec{\Sigma} \cdot \vec{V}), K\} = 0 \quad (6)$$

It is evident that the class of operators anticommuting with K (so-called K -odd operators) is much wider. Any operator of the form $\hat{O}(\vec{\Sigma} \cdot \vec{V})$, **where $\hat{O}$ commutes with K , but is otherwise arbitrary, also is a K -odd.** This fact will be used below.

Now one can proceed to the second stage of our problem: we wish to construct the K -odd operator A , that commutes with H . It is clear that there remains large freedom according to the above mentioned remark about $\hat{O}$ operators - one can take $\hat{O}$ into account or ignore it.

We have the following physically interesting vectors at hand which obey the requirements of our theorem. They are

$$\hat{\vec{r}} - \text{unit radius-vector and } \vec{p} - \text{linear momentum vector} \quad (7)$$

Both of them are perpendicular to $\vec{l}$ [14]. Thus, we choose the following K -odd terms

$$(\vec{\Sigma} \cdot \hat{\vec{r}}), \quad K(\vec{\Sigma} \cdot \vec{p}) \quad \text{and} \quad K\gamma^5 \quad (8)$$

and let probe the combination

$$A = x_1(\vec{\Sigma} \cdot \hat{\vec{r}}) + ix_2K(\vec{\Sigma} \cdot \vec{p}) + ix_3K\gamma^5f(r) \quad (9)$$

Here the coefficients $x_i (i = 1, 2, 3)$ are chosen in such a way that A operator is Hermitian for arbitrary real numbers and $f(r)$ is an arbitrary scalar function to be determined later from the symmetry requirements. Let's calculate

$$0 = [A, H] = (\vec{\Sigma} \cdot \hat{\vec{r}})\{x_2V'(r) - x_3f'(r)\} + 2i\beta K\gamma^5\left\{\frac{x_1}{r} - mx_3f(r)\right\} \quad (10)$$

We have a diagonal matrix in the first row, while the antidiagonal matrix in the second one. Therefore two equations follow:

$$x_2V'(r) = x_3f'(r), \quad x_3mf(r) = \frac{x_1}{r} \quad (11)$$

One can find from these equations:

$$V(r) = \frac{x_1}{x_2} \frac{1}{mr} \quad (12)$$

Therefore only the Coulomb potential corresponds to the above required $\pm\kappa$ degeneracy. The final form of obtained A operator is the following:

$$A = x_1 \left\{ (\vec{\Sigma} \cdot \hat{\vec{r}}) - \frac{i}{ma} K (\vec{\Sigma} \cdot \vec{p}) + \frac{i}{mr} K \gamma^5 \right\} \quad (13)$$

where unessential common factor x_1 may be omitted and after using known relations for Dirac matrices, this expression may be reduced to the form

$$A = \gamma^5 \left\{ \vec{\alpha} \cdot \hat{\vec{r}} - \frac{i}{ma} K \gamma^5 (H - \beta m) \right\} \quad (14)$$

Above and here a is a strength of Coulomb potential, $a = Z\alpha$. Precisely this operator is given in Johnson's and Lippmann's abstract [7].

What the real physical picture is standing behind this? Taking into account the relation

$$K (\vec{\Sigma} \cdot \vec{p}) = -i\beta \left(\vec{\Sigma} \cdot \frac{1}{2} [\vec{p} \times \vec{l} - \vec{l} \times \vec{p}] \right) \quad (15)$$

one can recast our operator in the following form

$$A = \vec{\Sigma} \cdot \left(\hat{\vec{r}} - \frac{i}{2ma} \beta [\vec{p} \times \vec{l} - \vec{l} \times \vec{p}] \right) + \frac{i}{mr} K \gamma^5$$

One can see that in the non-relativistic limit, i.e. $\beta \rightarrow 1$ and $\gamma^5 \rightarrow 0$ our operator reduces to

$$A \rightarrow A_{NR} = \vec{\sigma} \cdot \left(\hat{\vec{r}} - \frac{i}{2ma} [\vec{p} \times \vec{l} - \vec{l} \times \vec{p}] \right) \quad (16)$$

Note the LRL vector in the parenthesis of this equation. **Therefore relativistic supercharge reduces to the projection of the LRL vector on the electron spin direction. Precisely this operator was used in the case of Pauli electron [15].**

Because the Witten's Hamiltonian is

$$A^2 = 1 + \left(\frac{K}{a} \right)^2 \left(\frac{H^2}{m^2} - 1 \right) \quad (17)$$

and it consists only mutually commuting operators, it is possible their simultaneous diagonalization and replacement by corresponding eigenvalues. For instance, the energy of ground state is

$$E_0 = \left(1 - \frac{(Z\alpha)^2}{\kappa^2} \right)^{1/2} \quad (18)$$

By using the ladder procedure, familiar for SUSY quantum mechanics, the Sommerfeld formula can be easily derived [8].

Let's remark that if we include the Lorentz-scalar potential as well

$$H = \vec{\alpha} \cdot \vec{p} + \beta m + V(r) + \beta S(r) \quad (19)$$

this last Hamiltonian also commutes with K -operator, but *does not commute* with the above JL operator.

On the other hand, non-relativistic quantum mechanics is **indifferent** with regard of **the Lorentz variance properties** of potential. Therefore it is expected that in case of scalar

potential the description of hidden symmetry should also be possible. In other words, **the JL operator must be generalised** to this case.

For this purpose it is necessary to increase number of K -odd structures. One has to use our theorem in the part of additional $\hat{O}$ factors.

Now let us probe the following operator

$$B = x_1 (\vec{\Sigma} \cdot \hat{\vec{r}}) + x'_1 (\vec{\Sigma} \cdot \hat{\vec{r}}) H + ix_2 K (\vec{\Sigma} \cdot \vec{p}) + ix_3 K \gamma^5 f_1(r) + ix_4 K \gamma^5 \beta f_2(r) \quad (20)$$

We included $\hat{O} = H$ in the first structure and at the same time the matrix $\hat{O} = \beta$ in the third structure. Both of them commute with K and are permissible by our theorem. The form (20) is a minimal extension of the previous picture, because only the first order structures in $\hat{\vec{r}}$ and $\vec{p}$ participate. For turning to the previous case one must take $x_3 = x_4 = 0$ and $S(r)=0$. Calculation of relevant commutators give:

$$\begin{aligned} [B, H] = \gamma^5 \beta K \left\{ \frac{2ix_1}{r} - 2ix_3(m+S)f_1(r) + \frac{2ix'_1}{r} V(r) \right\} + \\ + K (\vec{\Sigma} \cdot \hat{\vec{r}}) \{ x_2 V'(r) - x_3 f'_1(r) \} + \\ + K \beta (\vec{\Sigma} \cdot \hat{\vec{r}}) \{ x_2 S'(r) - x_4 f'_2(r) \} + \\ + \gamma^5 K \left\{ \frac{2ix'_1(m+S)}{r} - 2ix_4(m+S)f_2(r) \right\} + \\ + \beta K \left\{ \frac{2ix'_1}{r} - 2ix_4 f_2(r) \right\} (\vec{\Sigma} \cdot \vec{p}) \end{aligned} \quad (21)$$

Equating this expression to zero, we derive matrix equation, then passing to 2×2 representation we must equate to zero the coefficients standing in fronts of diagonal and antidiagonal elements. In this way it follows equations:

(i) From antidiagonal structures (γ^5 , $\gamma^5 \beta K$)

$$\begin{aligned} \frac{x_1}{r} - x_3(m+S)f_1(r) + \frac{x'_1}{r} V(r) &= 0 \\ \frac{x'_1}{r}(m+S) - (m+S)f_2(r) &= 0 \end{aligned} \quad (22)$$

(ii) From diagonal structures ($K(\vec{\Sigma} \cdot \hat{\vec{r}})$, $K\beta(\vec{\Sigma} \cdot \hat{\vec{r}})$, $\beta K(\vec{\Sigma} \cdot \vec{p})$):

$$\begin{aligned} x_2 V'(r) - x_3 f'_1(r) &= 0 \\ \frac{x_1}{r} - x_2(m+S)V(r) + \frac{x'_1}{r} V(r) &= 0 \\ x_2 S'(r) - x_4 f'_2(r) &= 0 \\ \frac{x'_1}{r} - x_4 f_2(r) &= 0 \end{aligned} \quad (23)$$

Integrating the first and third equations in (23) for vanishing boundary conditions at infinity, we obtain

$$f_1(r) = \frac{x_2}{x_3} V(r), \quad f_2(r) = \frac{x_2}{x_4} S(r) \quad (24)$$

and taking into account the last equation from (23), we have

$$f_2(r) = \frac{x'_1}{x_4 r} \quad (25)$$

Therefore, according to (24), we obtain finally

$$S(r) = \frac{x'_1}{x_2 r} \quad (26)$$

So, the scalar potential must be Coulombic.

Inserting Eq.(24) into the first equation of (22) and solving for $V(r)$, one derives

$$V(r) = \frac{x_1}{r} \frac{1}{x_2(m + S) - \frac{x'_1}{r}} \quad (27)$$

At last, using here the expression (26), we find

$$V(r) = \frac{x_1}{x_2 m r} \quad (28)$$

Therefore we have asserted that the $\pm\kappa$ degeneracy is a symmetry of the Dirac equation only for Coulomb potential (for any general combination of Lorentz scalar and 4th component of a Lorentz vector).

Proceed from this fact, we believe that the nature of "accidental" or "hidden" symmetry of the Kepler problem is established - it is a $\pm\kappa$ degeneracy. Generators of this symmetry, A or B describe this degeneracy - they interchange these two values.

It is evident from an ordinary momentum algebra that the label $\kappa = j + 1/2$ takes place for aligned spin $j = l + 1/2$, i.e. for states $(s_{1/2}, p_{3/2}, \text{etc.})$, while $\kappa = -(j + 1/2)$ holds for unaligned spin or for states $(p_{1/2}, d_{3/2}, \text{etc.})$. **We see, that the $\pm\kappa$ degeneracy leads to the forbidden of the Lamb shift. Therefore absence of the Lamb shift in the Dirac equation for pure Coulomb potential is a consequence of the above mentioned symmetry, which in its deep meaning is related to the conservation of LRL vector (!).**

After all this by means of known ways the $SO(4)$ algebra could be constructed, manifestation of which in our macroworld is a conservation of the LRL vector and cosequently closeness of celestial orbits.

Therefore, it seems that the conservation of LRL vector is a macroscopic manifestation of symmetry, which is present in microworld.

Further, if we take into account above obtained solutions, one can reduce the B operator to more compact form

$$B = (\vec{\Sigma} \cdot \hat{r})(ma_V + a_S H) - \imath K \gamma^5 (H - \beta m) \quad (29)$$

where the following notations are used

$$a_V = -\frac{x_1}{x_2 m}, \quad a_S = -\frac{x'_1}{x_2}$$

Here a_i -s are the constants of corresponding Coulomb potentials

$$V(r) = -\frac{a_V}{r}, \quad S(r) = -\frac{a_S}{r}$$

In conclusion we want to remark, that this expression for B was mentioned recently by Leviatan [16], who used the radial decomposition and separation of spherical angles in the Dirac equation.

Our approach is 3-dimensional, without any referring to radial equation and, therefore is more general and rather easy. Moreover we find the source of mysterious "hidden symmetry of the Kepler problem, as the degeneracy on quantum level.

References

- [1] See e.g. R.C. O'Connell and K. Jagannathan. Am. J. Phys. **71**, 243 (2003);
- [2] W.Pauli. Z.Phys. **36**,336 (1926);
- [3] V.Fock. Z.Phys. **98**, 145 (1935);
- [4] J. Goldstein, C.Poole and J.Saiko. Classical Mechanics. 3rd Ed.(Pearson Education, 2003);
- [5] A.Bohm. Quantum Mechanics.(Springer-Verlag, New York,1993);
- [6] C. V. Sukumar, J.Phys.A: Math.Gen.**18**,L697 (1985);
- [7] M. H. Johnson and B.A. Lippman. Phys. Rev. **78**, 329(A), 1950.
- [8] H. Katsura, H. Aoki, J. Math. Phys. **47**, 032302 (2006);
- [9] T. T. Khachidze, A. A. Khelashvili, Mod. Phys. Lett.**A20**, 2277 (2005);
- [10] Proceedings - Fizika Visokikh Energii i Teoriya Elementarnikh Chastic, (Naukova Dumka, Kiev, 1967);
- [11] A. Sommerfeld. Atombau und Spectralinien (Fridr, Braunschweig, 1951), Vol.II;
- [12] E.Witten. Nucl.Phys. B188, 513 (1986);
- [13] T. T. Khachidze, A. A. Khelashvili, Am. J. Phys.**74**, 628, (2006); see also arXiv: quant-ph/0508122;
- [14] Constraints of this theorem are also satisfied by Laplace-Runge-Lenz vector $\vec{A} = \hat{r} - \frac{i}{2ma}[\vec{p} \times \vec{l} - \vec{l} \times \vec{p}]$, but this vector is associated to the Coulomb potential. Hence, we abstain from its consideration for now. Moreover $(\vec{\Sigma} \cdot \vec{A})$ is not an independent structure. It is expressible by two other structures, e.g. $(\vec{\Sigma} \cdot \vec{A}) = (\vec{\Sigma} \cdot \hat{r}) + \frac{i}{ma}\beta K(\vec{\Sigma} \cdot \vec{p})$;
- [15] R. G. Tangerman, J. A. Tjon. Phys. Rev.**A48**,1089 (1998);
- [16] A.Leviatan, Phys.Rev.Lett.**92**, 202501 (2004).